## Asian American, Native Hawaiian, & Pacific Islander Health Inequities Compared to Non-Hispanic Whites

Racial and ethnic health inequities are undermining our communities and our entire health care system. Asian Americans (AAs) and Native Hawaiians and Pacific Islanders (NHPIs) experience significant health inequities that are often inadequately reported or not reported at all. AAs and NHPIs are the fastest growing racial groups in the nation, and one of the most diverse, tracing their heritage to more than 50 different countries. Yet data on the AA and NHPI population is lumped together, masking the distinct health needs within AA and NHPI whites with that of AA and NHPIs, with disaggregated data where available.

#### ASIAN AMERICAN, NATIVE HAWAIIAN, & PACIFIC ISLANDER HEALTH INEQUITIES: ADULTS

#### Stomach Cancer

Women

76%

more likely to develop stomach cancer (Asian and Pacific Islanders)2

as likely to die from stomach cancer (Asian and Pacific Islanders)3

Men

more likely to develop stomach cancer (Asian and Pacific Islanders)<sup>2</sup>

as likely to die from stomach cancer (Asian and Pacific Islanders)3

## **Hepatitis**

**68**%

more likely to contract hepatitis A (Asian and Pacific Islanders)5

x18as likely to contract

chronic hepatitis B (Asian and Pacific Islanders)5

## **Diabetes**

**76**%

diabetic (Native

diabetic (Asian-Indian Americans)7

more likely to develop

(Asian Americans)8

## **Obesity**

more likely to be obese (Native Hawaiian and other Pacific Islanders)6



# Pre-natal

more likely to receive late or no prenatal care (Chinese Americans)1

as likely to receive late or no prenatal care (Hawaiian and part Hawaiian)1

### Tuberculosis\*

**x5** 

tuberculosis (Asian Americans)4

as likely to contract

**x16** as likely to contract

Hawaiian and Pacific Islanders)4 \*Among U.S.-born persons

tuberculosis (Native

more likely to be

Hawaiian and Pacific Islanders)6

more likely to be

50%

end-stage renal disease

### Liver Cancer Women

more likely to develop

liver and IBD cancer (Asian and Pacific Islanders)2

more likely to die from

\*IBD = Intrahepatic Bile

47%

liver and IBD cancer (Asian and Pacific

Islanders)3

**Duct Cancer** 

66%

Men

more likely to develop liver and IBD cancer

(Asian and Pacific Islanders)2

**58**% more likely to die from

**Duct Cancer** 

liver and IBD cancer (Asian and Pacific Islanders)3 \*IBD = Intrahepatic Bile

Our common prosperity demands a health system where everyone can attain the best possible health and health care. The roots of health and ensure a better future for all.





AT FAMILIESUSA



NOTE: Data includes Asian Americans, Native Hawaiians and Pacific Islanders, unless otherwise stated.

<sup>1</sup> Child Trends DataBank, Late or No Prenatal Care: Indicators of Child and Youth Well-Being, 2015 <sup>2</sup> NIH, Cancer Stat Facts

- <sup>3</sup> CDC, National Vital Statistics Reports, Deaths: Final Data for 2016
- <sup>4</sup> CDC, Morbidity and Mortality Weekly Report, Tuberculosis United States, 2017
- <sup>5</sup> CDC, Morbidity and Mortality Weekly Report, Summary of Notifiable Infectious Diseases and
- Conditions United States, 2015 6 CDC, Summary Health Statistics: National Health Interview Survey, 2016
- <sup>7</sup> CDC, National Diabetes Statistics Report, 2017
- <sup>8</sup> United States Renal Data System, 2015 USRDS Annual Data Report

\*Families USA thanks the Asian & Pacific Islander American Health Forum for their assistance in preparing this piece

January 2019 FamiliesUSA.org